

GEOMETRIC CONSTRUCTION OF GENERATORS OF COHA OF DOUBLED QUIVER

ZONGBIN CHEN

ABSTRACT. Let Q be the double of a quiver. According to Efimov, Kontsevich and Soibelman, the cohomological Hall algebra (CoHA) associated to Q is a free super-commutative algebra. In this short note, we confirm a conjecture of Hausel which gives a geometric realisation of the generators of the CoHA.

1. INTRODUCTION

Let $Q = (I, \Omega)$ be a quiver with a set of vertices I and with a_{ij} arrows from $i \in I$ to $j \in I$. For each dimension vector $\gamma = (\gamma^i)_{i \in I} \in \mathbf{Z}_{\geq 0}^I$, we have the affine $\mathbf{C}$ -variety M_γ of representations of Q in complex coordinate space $\bigoplus_{i \in I} \mathbf{C}^{\gamma^i}$. It is acted on by the complex algebraic group $G_\gamma = \prod_{i \in I} \mathrm{GL}_{\gamma^i}(\mathbf{C})$ and the action factors through $PG_\gamma := G_\gamma/\mathbb{G}_m$, where $\mathbb{G}_m$ is embedded diagonally in G_γ . We denote by $[M_\gamma/G_\gamma]$ the moduli stack of representations of Q of dimension γ .

For $\gamma_1, \gamma_2 \in \mathbf{Z}_{\geq 0}^I$, let M_{γ_1, γ_2} be the subvariety of $M_{\gamma_1 + \gamma_2}$ consisting of the representations of Q such that the subspace $\bigoplus_{i \in I} \mathbf{C}^{\gamma_1^i} \subset \bigoplus_{i \in I} (\mathbf{C}^{\gamma_1^i} \oplus \mathbf{C}^{\gamma_2^i})$ forms a sub representation. Let $G_{\gamma_1, \gamma_2} \subset G_{\gamma_1 + \gamma_2}$ be the subgroup preserving $\bigoplus_{i \in I} \mathbf{C}^{\gamma_1^i}$. Then $[M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}]$ is the moduli stack classifying the extensions of representations of Q of dimension vector γ_2 by that of dimension vector γ_1 . We have the correspondence

$$\begin{array}{ccc} & [M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}] & \\ p \swarrow & & \searrow q \\ [M_{\gamma_1}/G_{\gamma_1}] \times [M_{\gamma_2}/G_{\gamma_2}] & & [M_{\gamma_1 + \gamma_2}/G_{\gamma_1 + \gamma_2}], \end{array}$$

from which Kontsevich and Soibelman [KS] have constructed an associative algebra structure on

$$\mathcal{H} := \bigoplus_{\gamma \in \mathbf{Z}_{\geq 0}^I} \mathcal{H}_\gamma, \quad \mathcal{H}_\gamma := H^*([M_\gamma/G_\gamma]) = H_{G_\gamma}^*(M_\gamma),$$

which is called the *cohomological Hall algebra* (CoHA) of the quiver Q . Here and after, all the cohomological groups take coefficients in $\mathbf{Q}$. The resulting product has a shift in cohomological degree:

$$H_{G_{\gamma_1}}^*(M_{\gamma_1}) \times H_{G_{\gamma_2}}^*(M_{\gamma_2}) \rightarrow H_{G_{\gamma_1 + \gamma_2}}^{*-2\chi_Q(\gamma_1, \gamma_2)}(M_{\gamma_1 + \gamma_2}),$$

where

$$\chi_Q(\gamma_1, \gamma_2) = \sum_{i \in I} \gamma_1^i \gamma_2^i - \sum_{i, j \in I} a_{i, j} \gamma_1^i \gamma_2^j.$$

Suppose that the quiver Q is symmetric, i.e. $a_{ij} = a_{ji}$. In this case, $\mathcal{H}$ has more structures. First of all, one can make $\mathcal{H}$ into a $(\mathbf{Z}_{\geq 0}^I, \mathbf{Z})$ -graded algebra, by requiring elements in $H_{G_\gamma}^k(M_\gamma)$ to be of bidegree $(\gamma, k + \chi_Q(\gamma, \gamma))$. Secondly, following Efimov [E], we can twist the multiplication by a sign such that $(\mathcal{H}, *)$ is a super-commutative algebra with respect to the $\mathbf{Z}$ -grading. In fact, for $a_{\gamma, k} \in \mathcal{H}_{\gamma, k}$, $a_{\gamma', k'} \in \mathcal{H}_{\gamma', k'}$, we have

$$a_{\gamma, k} a_{\gamma', k'} = (-1)^{\chi_Q(\gamma, \gamma')} a_{\gamma', k'} a_{\gamma, k}.$$

We can find a bilinear form $\psi : (\mathbf{Z}/2)^I \times (\mathbf{Z}/2)^I \rightarrow \mathbf{Z}/2$ such that

$$\psi(\gamma_1, \gamma_2) + \psi(\gamma_2, \gamma_1) \equiv \chi_Q(\gamma_1, \gamma_2) + \chi_Q(\gamma_1, \gamma_1) \chi_Q(\gamma_2, \gamma_2) \pmod{2}.$$

Then the twisted product on $\mathcal{H}$ is defined to be

$$a_{\gamma, k} * a_{\gamma', k'} = (-1)^{\psi(\bar{\gamma}, \bar{\gamma}')} a_{\gamma, k} \cdot a_{\gamma', k'},$$

where $\bar{\gamma}$ is the image of γ in $(\mathbf{Z}/2)^I$.

For the symmetric quiver Q , it is conjectured by Kontsevich and Soibelman [KS] and proved by Efimov [E], that the $(\mathbf{Z}_{\geq 0}^I, \mathbf{Z})$ -graded algebra $(\mathcal{H}, *)$ is a free super-commutative algebra generated by a $(\mathbf{Z}_{\geq 0}^I, \mathbf{Z})$ -graded vector space V of the form $V = V^{\text{prim}} \otimes \mathbf{Q}[x]$, with x an element of degree $(0, 2)$, and for all $\gamma \in \mathbf{Z}_{\geq 0}^I$ the vector space $V_{\gamma, k}^{\text{prim}}$ is non-zero for only finitely many k . Geometrically, via the isogeny $G_\gamma \rightarrow PG_\gamma \times \mathbb{G}_m$, we have

$$\mathcal{H}_\gamma = H_{G_\gamma}^*(M_\gamma) \cong H_{PG_\gamma}^*(M_\gamma) \otimes H_{\mathbb{G}_m}^*(\text{pt}),$$

and it gives the above factorisation $V = V^{\text{prim}} \otimes \mathbf{Q}[x]$. Let M_γ^{st} be the stable part of M_γ with respect to the 0-stability condition, i.e. M_γ^{st} consists of all the simple Q -modules. Then as we will see in theorem 2.5, $V_{\gamma, * + \chi_Q(\gamma, \gamma)}^{\text{prim}}$ is in fact the pure part of $H^*([M_\gamma^{st}/PG_\gamma])$, where the word “pure” refers to the mixed Hodge structure on the cohomological group. Let $c_{\gamma, k} = \dim_{\mathbf{C}} V_{\gamma, k}^{\text{prim}}$, the above result implies that the quantum Donaldson-Thomas invariants of the quiver Q without potential and with 0-stability condition is

$$\Omega(\gamma)(q) = \sum_{k \in \mathbf{Z}} c_{\gamma, k} q^{k/2} \in \mathbf{Z}[q^{\pm \frac{1}{2}}].$$

In particular, the coefficients are positive.

On the other hand, in the work of Hausel, Letellier and Rodriguez-Villegas [HLR], they found another expression for the quantum Donaldson-Thomas invariants. From now on, we work with quivers $Q = (I, \Omega)$ that are the double of another quiver, i.e. $\Omega = \Omega_0 \sqcup \Omega_0^{\text{op}}$, where Ω_0^{op} is obtained by reversing all the arrows in Ω_0 . In this case, M_γ is endowed with a G_γ -invariant holomorphic symplectic form ω . Let $\mu : M_\gamma \rightarrow \mathfrak{g}_\gamma^0$ be the corresponding moment map, here $\mathfrak{g}_\gamma^0$ is the trace 0 part of $\mathfrak{g}_\gamma := \text{Lie}(G_\gamma)$. Let $\mathcal{O}$ be the G_γ -orbit of a *generic* (to be explained below) regular semisimple element in $\mathfrak{g}_\gamma^0$. The group PG_γ acts freely on $\mu^{-1}(\mathcal{O})$ and we have the geometric quotient $\mu^{-1}(\mathcal{O})/PG_\gamma$, which is a smooth

quasi-projective algebraic variety. Furthermore, the Weyl group W_γ of G_γ acts on the cohomological groups $H^*(\mu^{-1}(\mathcal{O})/PG_\gamma)$. One of the main results of [HLR] states that

$$\Omega(\gamma)(q) = q^{\frac{1}{2}\chi_Q(\gamma,\gamma)} \sum_i \dim(H^{2i}(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}) q^i.$$

Based on this result, Hausel conjectured that the cohomological groups

$$H^k(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}$$

are geometric realisations of the generating set $V_{\gamma,k+\chi_Q(\gamma,\gamma)}^{\text{prim}}$. (Of course, the conjecture is meaningful only when $\Omega(\gamma)(q)$ is non-zero. We will always impose this condition). In this article, we confirm this conjecture. Our construction goes as follows:

Let $\chi : \mathfrak{g}_\gamma^0 \rightarrow \mathfrak{g}_\gamma^0 // G_\gamma \cong \mathfrak{t}_\gamma^0 // W_\gamma$ be the characteristic morphism. Following Ginzburg [G], we consider the composition $f : M_\gamma \xrightarrow{\mu} \mathfrak{g}_\gamma^0 \xrightarrow{\chi} \mathfrak{t}_\gamma^0 // W_\gamma$. Let

$$\mathfrak{t}_\gamma^{0,\text{gen}} := \mathfrak{t}_\gamma^{0,\text{reg}} \setminus \bigcup_{\substack{\gamma_1, \gamma_2 \in \mathbf{Z}_{\geq 0}^I, \\ \gamma_1 + \gamma_2 = \gamma}} W_\gamma \cdot (\mathfrak{t}_{\gamma_1}^0 \oplus \mathfrak{t}_{\gamma_2}^0),$$

we call conjugates of elements in it *generic* regular semisimple elements. Let

$$U_\gamma := f^{-1}(\mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma),$$

then the group PG_γ acts freely on U_γ and the quotient U_γ/PG_γ is a quasi-projective algebraic variety. Furthermore, the restriction of the morphism f to U_γ descends to a morphism $\bar{f} : U_\gamma/PG_\gamma \rightarrow \mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$. We'll prove that it makes U_γ/PG_γ a fiber bundle on $\mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$ with fibers isomorphic to $\mu^{-1}(\mathcal{O})/PG_\gamma$. Our main result is the following:

Theorem 1.1. *The pure part of $H^*(U_\gamma/PG_\gamma)$ is equal to $H^*(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}$. The restriction $H_{PG_\gamma}^*(M_\gamma) \rightarrow H^*(U_\gamma/PG_\gamma)$ factors through and is surjective onto the pure part of the latter, and its restriction to $V_{\gamma,*+\chi_Q(\gamma,\gamma)}^{\text{prim}}$ induces an isomorphism*

$$V_{\gamma,*+\chi_Q(\gamma,\gamma)}^{\text{prim}} \cong H^*(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}.$$

2. PROOF OF THE MAIN THEOREM

We begin by recalling the construction of Kontsevich and Soibelman of the cohomological Hall algebra. Given two vectors $\gamma_1, \gamma_2 \in \mathbf{Z}_{\geq 0}^I$, let $\gamma = \gamma_1 + \gamma_2$. The product $\mathcal{H}_{\gamma_1} \times \mathcal{H}_{\gamma_2} \rightarrow \mathcal{H}_\gamma$ is defined to be the composition of the Künneth isomorphism

$$H_{G_{\gamma_1}}^*(M_{\gamma_1}) \otimes H_{G_{\gamma_2}}^*(M_{\gamma_2}) \cong H_{G_{\gamma_1} \times G_{\gamma_2}}^*(M_{\gamma_1} \times M_{\gamma_2}),$$

and the following morphisms

$$(1) \quad H_{G_{\gamma_1} \times G_{\gamma_2}}^*(M_{\gamma_1} \times M_{\gamma_2}) \cong H_{G_{\gamma_1, \gamma_2}}^*(M_{\gamma_1, \gamma_2}) \xrightarrow{\phi_1} H_{G_{\gamma_1, \gamma_2}}^{*+2c_1}(M_\gamma) \xrightarrow{\phi_2} H_{G_\gamma}^{*+2c_1+2c_2}(M_\gamma),$$

where $c_1 = \dim_{\mathbf{C}} M_\gamma - \dim_{\mathbf{C}} M_{\gamma_1, \gamma_2}$ and $c_2 = \dim_{\mathbf{C}} G_{\gamma_1, \gamma_2} - \dim_{\mathbf{C}} G_\gamma$, and the first isomorphism is induced by the fibrations in affine spaces

$$M_{\gamma_1, \gamma_2} \rightarrow M_{\gamma_1} \times M_{\gamma_2}, \quad G_{\gamma_1, \gamma_2} \rightarrow G_{\gamma_1} \times G_{\gamma_2},$$

and the other morphisms ϕ_1, ϕ_2 are natural push forwards.

Lemma 2.1. *Under the restriction $H^*([M_\gamma/G_\gamma]) \rightarrow H^*([M_\gamma^{st}/G_\gamma])$, the image of*

$$\bigoplus_{\substack{\gamma_1, \gamma_2 \in \mathbf{Z}_{\geq 0}^I \\ \gamma_1 + \gamma_2 = \gamma}} \mathcal{H}_{\gamma_1} \times \mathcal{H}_{\gamma_2}$$

in $\mathcal{H}_\gamma = H^*([M_\gamma/G_\gamma])$ goes to 0.

Proof. By the definition of Gysin map, the morphism ϕ_1 in the composition (1) factorises as

$$H^*([M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}]) \rightarrow H_{[M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}]}^{*+2c_1}([M_\gamma/G_{\gamma_1, \gamma_2}]) \rightarrow H^{*+2c_1}([M_\gamma/G_{\gamma_1, \gamma_2}]).$$

Using the long exact sequence

$$\cdots \rightarrow H_{[M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}]}^*([M_\gamma/G_{\gamma_1, \gamma_2}]) \rightarrow H^*([M_\gamma/G_{\gamma_1, \gamma_2}]) \rightarrow H^*([M_\gamma/G_{\gamma_1, \gamma_2}] \setminus [M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}]) \rightarrow \cdots,$$

we see that $\text{Im}(\phi_1)$ goes to 0 when we restrict it to

$$H^{*+2c_1}([M_\gamma/G_{\gamma_1, \gamma_2}] \setminus [M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}]).$$

Since $[M_\gamma^{st}/G_{\gamma_1, \gamma_2}]$ is contained in $[M_\gamma/G_{\gamma_1, \gamma_2}] \setminus [M_{\gamma_1, \gamma_2}/G_{\gamma_1, \gamma_2}]$, $\text{Im}(\phi_1)$ vanishes when we restrict it further to

$$H^{*+2c_1}([M_\gamma^{st}/G_{\gamma_1, \gamma_2}]).$$

Now applying ϕ_2 , we see that $\text{Im}(\phi_2 \circ \phi_1)$ vanishes when we restrict it to $H^{*+2c_1+2c_2}([M_\gamma^{st}/G_\gamma])$. $\square$

We need some preliminary results before proceeding to the proof of the main theorem. Given an element $t = (t_i)_{i \in I} \in \mathfrak{t}_\gamma^{0, \text{gen}}$, let $\mathcal{O}$ be its orbit under the action of PG_γ by conjugation. Recall that Crawley-Boevey [CB2] has identified the geometric quotient $\mu^{-1}(\mathcal{O})/PG_\gamma$ with a quiver variety: Let $\tilde{Q}$ be the quiver obtained from Q by attaching to each vertex $i \in I$ a leg of length $\gamma_i - 1$. More precisely, vertices of $\tilde{Q}$ are labeled $[i, j]$, $i \in I, j = 0, \dots, \gamma_i - 1$, and we identify $[i, 0]$ with i . Besides the arrows in Q , the new arrows in $\tilde{Q}$ are $[i, j] \rightrightarrows [i, j+1]$ for each $i \in I, j = 0, \dots, \gamma_i - 2$. The new dimension vector of $\tilde{Q}$ is defined to be $\tilde{\gamma}_{[i, j]} = \gamma_i - j$. Again, we have the moment map $\tilde{\mu} : M_{\tilde{\gamma}}^{\tilde{Q}} \rightarrow \mathfrak{g}_{\tilde{\gamma}}^0$, where $M_{\tilde{\gamma}}^{\tilde{Q}}$ is the space of representations of $\tilde{Q}$ of dimension vector $\tilde{\gamma}$. For each $i \in I$, let $t_{i,1}, \dots, t_{i,\gamma_i}$ be the eigenvalues of t_i . Define $\lambda = (\lambda_{[i, j]}) \in \mathfrak{g}_{\tilde{\gamma}}^0$ to be

$$\begin{aligned} \lambda_{[i, 0]} &= -t_{i,1}, \\ \lambda_{[i, j]} &= t_{i,j} - t_{i,j+1}, \quad j = 1, \dots, \gamma_i - 1. \end{aligned}$$

Notice that $\tilde{\gamma} \cdot \lambda = 0$. Now the result of Crawley-Boevey [CB2] states that

$$(2) \quad \mu^{-1}(\mathcal{O})/PG_\gamma \cong \tilde{\mu}^{-1}(\lambda)/PG_{\tilde{\gamma}}.$$

Moreover, according to [HLR], corollary 1.6 (iv), $\Omega(\gamma)(q)$ is non-zero if and only if $\tilde{\gamma}$ is a positive root of Q' , here we write $\tilde{Q}$ as the double of another quiver Q' .

Lemma 2.2. *The morphism $\bar{f} : U_\gamma/PG_\gamma \rightarrow \mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$ makes U_γ/PG_γ a fiber bundle over $\mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$. Moreover, the sheaf $R^i \bar{f}_* \mathbf{Q}$ is constant on the étale neighbourhood $\mathfrak{t}_\gamma^{0,\text{gen}} \rightarrow \mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$ of $\mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$.*

Proof. This is basically the lemma 48 of [M], with one difference. As in the proof of [HLR] theorem 2.3, Maffei works with quiver without loops, but his proof carries over in our case. In his proof, the important point is the surjectivity of $\bar{f}$ (or rather the hyperkähler moment map on the generic locus, but this can be reduced to $\bar{f}$ by hyperkähler rotation). According to [CB1], theorem 4.4, this is fulfilled in our situation since $\mu^{-1}(\mathcal{O})/PG_\gamma \cong \tilde{\mu}^{-1}(\lambda)/PG_{\tilde{\gamma}}$ and $\tilde{\gamma} \cdot \lambda = 0$, taking into account that $\tilde{\gamma}$ is a positive root of Q' . $\square$

Lemma 2.3 (Crawley-Boevey-van den Bergh). *The smooth quasi-projective algebraic variety $\mu^{-1}(\mathcal{O})/PG_\gamma$ has pure mixed Hodge structure.*

Proof. This is a corollary of [CBB], §2.4, taking into account the isomorphism (2). $\square$

Now we can prove the first part of theorem 1.1.

Theorem 2.4. *The pure part of $H^*(U_\gamma/PG_\gamma)$ is equal to $H^*(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}$.*

Proof. Consider the fiber bundle $\bar{f} : U_\gamma/PG_\gamma \rightarrow \mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$. By lemma 2.2, the sheaf $R^i \bar{f}_* \mathbf{Q}$ is constant on the étale neighbourhood $\mathfrak{t}_\gamma^{0,\text{gen}} \rightarrow \mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$ of $\mathfrak{t}_\gamma^{0,\text{gen}}/W_\gamma$, so we get the Hochschild-Serre spectral sequence,

$$E_2^{p,q} = H^p(W_\gamma, H^q(\mathfrak{t}_\gamma^{0,\text{gen}}, R\bar{f}_* \mathbf{Q})) \implies H^{p+q}(U_\gamma/PG_\gamma).$$

Since W_γ is a finite group, we have $E_2^{p,q} = 0$ for $p \neq 0$. So the spectral sequence degenerates, and we get

$$\begin{aligned} H^q(U_\gamma/PG_\gamma) &= (H^q(\mathfrak{t}_\gamma^{0,\text{gen}}, R\bar{f}_* \mathbf{Q}))^{W_\gamma} \\ &= \left(\bigoplus_{q_1+q_2=q} H^{q_1}(\mathfrak{t}_\gamma^{0,\text{gen}}) \otimes H^{q_2}(\mu^{-1}(\mathcal{O})/PG_\gamma) \right)^{W_\gamma}. \end{aligned}$$

Since $\mathfrak{t}_\gamma^{0,\text{gen}}$ is the complement of unions of sufficiently many hyperplanes in the vector space $\mathfrak{t}_\gamma^0$, one proves easily by induction on the number of hyperplanes that the mixed Hodge structure of $H^i(\mathfrak{t}_\gamma^{0,\text{gen}})$ is not pure if $i \neq 0$. So the pure part of $H^q(U_\gamma/PG_\gamma)$ is exactly $H^q(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}$. $\square$

To prove the second result in the main theorem, we need some facts from algebraic stacks. We refer the reader to Olsson-Laszlo [OL 1] and Sun [S] for the proofs. Although they work over the finite fields, their results apply in our situation since the quiver varieties are in fact $\mathbf{Z}$ -schemes. The moduli stack $[M_\gamma/PG_\gamma]$ is a smooth Artin stack over $\mathbf{C}$ with dimension

$d_\gamma = \dim(M_\gamma) - \dim(PG_\gamma)$. It has dualizing complex $\mathbf{Q}(d_\gamma)[2d_\gamma]$, and we have the Poincaré duality, which is a perfect non-degenerate bilinear pairing

$$H^i([M_\gamma/PG_\gamma]) \times H_c^{2d_\gamma-i}([M_\gamma/PG_\gamma]) \rightarrow \mathbf{Q}(d_\gamma).$$

Using the fibration $[M_\gamma/PG_\gamma] \rightarrow [\text{pt}/PG_\gamma]$, we have $H^i([M_\gamma/PG_\gamma]) = H_{PG_\gamma}^i(\text{pt})$ is pure of weight i , the groups $H_c^{2d_\gamma-i}([M_\gamma/PG_\gamma])$ are all pure of weight $2d_\gamma - i$.

Furthermore, let $Z_\gamma = M_\gamma \setminus U_\gamma$, then $H_c^i([Z_\gamma/PG_\gamma])$ is of weight less than or equal to i . This is essentially [Weil II]. More precisely, as in [BL], let $\{E_n \rightarrow B_n\}_{n \in \mathbf{N}}$ be an injective system of finite dimensional n -acyclic approximation to the universal PG_γ -torsor $E \rightarrow B$, then

$$H_c^i([Z_\gamma/PG_\gamma]) = \lim_{n \rightarrow \infty} H_c^{i+2\dim(E_n)}(Z_\gamma \times_{PG_\gamma} E_n)(-\dim(E_n)).$$

Now it suffices to apply [Weil II] to the right hand side.

Proof of the second part of theorem 1.1. We have the long exact sequence

$$(3) \quad \cdots \rightarrow H_c^{i-1}([Z_\gamma/PG_\gamma]) \rightarrow H_c^i([U_\gamma/PG_\gamma]) \rightarrow H_c^i([M_\gamma/PG_\gamma]) \rightarrow H_c^i([Z_\gamma/PG_\gamma]) \rightarrow \cdots.$$

Since $H_c^{i-1}([Z_\gamma/PG_\gamma])$ is of weight less than or equal to $i-1$, the pure part of $H_c^i([U_\gamma/PG_\gamma])$ injects into $H_c^i([M_\gamma/PG_\gamma])$. Taking Poincaré duality, we have that $H^{2d_\gamma-i}([M_\gamma/PG_\gamma])$ maps onto the pure part of $H^{2d_\gamma-i}([U_\gamma/PG_\gamma])$. By theorem 2.4, we have surjective morphism

$$H^j([M_\gamma/PG_\gamma]) \twoheadrightarrow H^j(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}, \quad \forall j.$$

By the definition of $\mathfrak{t}_\gamma^{0,\text{gen}}$, we find easily that $U_\gamma \subset M_\gamma^{st}$. So the above map factorize by

$$(4) \quad H^j([M_\gamma/PG_\gamma]) \rightarrow H^j(M_\gamma^{st}/PG_\gamma) \twoheadrightarrow H^j(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}.$$

By lemma 2.1, the first arrow has the same image as its restriction to $V_{\gamma,j+\chi_Q(\gamma,\gamma)}^{\text{prim}}$, so we get a surjective morphism

$$V_{\gamma,j+\chi_Q(\gamma,\gamma)}^{\text{prim}} \twoheadrightarrow H^j(\mu^{-1}(\mathcal{O})/PG_\gamma)^{W_\gamma}.$$

By the result of [HLR] recalled in the introduction, they have the same dimension, so they are isomorphic. □

Similar arguments can be used to show the following variant of the geometric construction.

Theorem 2.5. *The restriction*

$$H^*([M_\gamma/PG_\gamma]) \rightarrow H^*([M_\gamma^{st}/PG_\gamma])$$

induces an isomorphism

$$V_{\gamma,*+\chi_Q(\gamma,\gamma)}^{\text{prim}} \cong PH^*([M_\gamma^{st}/PG_\gamma])$$

where $PH^*([M_\gamma^{st}/PG_\gamma])$ is the pure part of $H^*([M_\gamma^{st}/PG_\gamma])$.

Proof. The proof is almost the same as that of the main theorem, we indicate only the differences. Using an exact sequence as (3), with the pair (U_γ, Z_γ) replaced by $(M_\gamma^{st}, M_\gamma \setminus M_\gamma^{st})$, we can show that the restriction $H^*([M_\gamma/P G_\gamma]) \rightarrow H^*([M_\gamma^{st}/P G_\gamma])$ factors through and is surjective onto $PH^*([M_\gamma^{st}/P G_\gamma])$. Again by lemma 2.1, we get the surjection

$$V_{\gamma, * + \chi_Q(\gamma, \gamma)}^{\text{prim}} \twoheadrightarrow PH^*([M_\gamma^{st}/P G_\gamma]).$$

Now observe that the second morphism in the factorisation (4) has the same image as that of

$$PH^j([M_\gamma^{st}/P G_\gamma]) \rightarrow H^j(\mu^{-1}(\mathcal{O})/P G_\gamma)^{W_\gamma},$$

since the morphism preserves the weights of the cohomological groups and $H^j(\mu^{-1}(\mathcal{O})/P G_\gamma)^{W_\gamma}$ is pure of weight j . So the factorisation (4) becomes

$$(5) \quad V_{\gamma, j + \chi_Q(\gamma, \gamma)}^{\text{prim}} \twoheadrightarrow PH^j([M_\gamma^{st}/P G_\gamma]) \twoheadrightarrow H^j(\mu^{-1}(\mathcal{O})/P G_\gamma)^{W_\gamma}.$$

Now that the composition is an isomorphism by our main theorem, all the arrows in (5) are isomorphisms. □

ACKNOWLEDGEMENTS

We are very grateful to Tamás Hausel for having explained to us his conjecture, and to Ben Davison for pointing out a bug in the proof. We also want to thank an anonymous referee for his careful readings and helpful suggestions.

REFERENCES

- [BL] J. Bernstein, V. Lunts, *Equivariant sheaves and functors*. Lecture Notes in Mathematics 1578. Springer-Verlag, Berlin, 1994.
- [CB1] W. Crawley-Boevey, *Geometry of the moment map for representations of quivers*. Compositio Math. 126 (2001), no. 3, 257–293.
- [CB2] W. Crawley-Boevey, *On matrices in prescribed conjugacy classes with no common invariant subspace and sum zero*, Duke Math. J. 118 (2003), no. 2, 339–352.
- [CBB] W. Crawley-Boevey, M. van den Bergh, *Absolutely indecomposable representations and Kac-Moody Lie algebras*, Invent. Math. 155 (2004), no. 3, 537–559.
- [E] A. I. Efimov, *Cohomological Hall algebra of a symmetric quiver*. Compos. Math. 148 (2012), no. 4, 1133–1146.
- [HLR] T. Hausel, E. Letellier, F. Rodriguez-Villegas, *Positivity for Kac polynomials and DT-invariants of quivers*. Ann. of Math. (2) 177 (2013), no. 3, 1147–1168.
- [G] V. Ginzburg, *Talk at Luminy*, 2012.
- [KS] M. Kontsevich, Y. Soibelman, *Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson-Thomas invariants*. Commun. Number Theory Phys. 5 (2011), no. 2, 231–352.
- [M] A. Maffei, *A remark on quiver varieties and Weyl groups*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 1 (2002), no. 3, 649–686.
- [OL 1] Y. Laszlo, M. Olsson, *The six operations for sheaves on Artin stacks. I. Finite coefficients*. Publ. Math. Inst. Hautes Études Sci. No. 107 (2008), 109–168.

- [OL 2] Y. Laszlo, M. Olsson, *The six operations for sheaves on Artin stacks. II. Adic coefficients*. Publ. Math. Inst. Hautes Études Sci. No. 107 (2008), 169–210.
- [S] S. Sun, *Decomposition theorem for perverse sheaves on Artin stacks over finite fields*. Duke Math. J. 161 (2012), no. 12, 2297–2310.
- [Weil II] P. Deligne, *La conjecture de Weil, II*, Publ. Math. IHES, 52, 1980, 137–252.

EPFL SB MATHGEOM/GEOM, MA B1 447, LAUSANNE, CH-1015, SWITZERLAND
E-mail address: `zongbin.chen@epfl.ch`